

PF-0049P

## A NOVEL HUMAN JAK2 KINASE

## FIELD OF THE INVENTION

The present invention relates to a novel, human Jak2 kinase isolated from human placenta and to the use of this novel protein and its nucleic acid sequence in the diagnosis, study, <sup>prevention</sup>~~prevention~~ and treatment of disease.

## BACKGROUND OF THE INVENTION

JAK kinases are Janus family nonreceptor protein-tyrosine kinases (NR-PTK) that lack transmembrane regions and form functional complexes with the intercellular regions of other cell surface receptors. They were first identified as the products of mutant oncogenes in cancer cells where their activation was no longer subject to <sup>Jak3</sup>~~normal~~ cellular controls. Described JAK kinases include Jak1, Jak2, and ~~Jak3~~ <sup>addition</sup> which all share the conserved kinase domain. In ~~addition~~ these proteins have 5 to 100 amino acid residues located on either side of, or inserted into loops of, the carboxyterminal kinase domain which allow the regulation of each kinase as it recognizes and interacts with its target protein. Known target proteins include growth hormone receptor, prolactin receptor, erythropoietin receptor, cytokine receptors and others which utilize the common chain known as gp130. These receptors are unique both in their ability to recruit multiple PTKs and in the diversity of their responses within different cell types (Taniguchi T (1995) Science 268:251-55). Genetic evidence places these kinases in the interferon  $\alpha$  and  $\gamma$  signal transduction pathways which are widely expressed in mammalian cells.

Kinases regulate many different cell proliferation, differentiation, and signaling processes by adding phosphate groups to proteins. The high energy phosphate which drives activation is generally transferred from adenosine triphosphate molecules (ATP) to a particular protein by the PTKs, and the transfer process is roughly analogous to turning on a molecular switch. When the switch goes on, the kinase activates a metabolic enzyme, regulatory protein, receptor, cytoskeletal protein, ion channel or pump, transcription factor, or another kinase. For example, in their normal role, the JAK NR-PTKs are capable of regulating tyrosine phosphorylation of STAT proteins, signal transducers and activators of transcription, such that they translocate to the nucleus and bind DNA (David M et al. (1995)

1 Science 269:1721-1723). In contrast, uncontrolled kinase signaling has  
 been implicated in inflammation, oncogenesis, arteriosclerosis, and  
 psoriasis.

6 Almost all kinases contain a similar 250-300 amino acid catalytic  
 domain. The N-terminal domain generally folds into a two-lobed structure  
 to bind and orient ATP (or GTP) donor molecules. The larger C terminal  
 lobe binds the protein substrate and carries out the transfer of phosphate  
 from ATP to the hydroxyl group of a tyrosine residue. The primary  
 structure of the kinase domain is conserved in the residues: G<sub>50</sub> and G<sub>52</sub> in  
 subdomain I, K<sub>72</sub> in subdomain II, G<sub>91</sub> in subdomain III, E<sub>208</sub> in subdomain  
 11 VIII, D<sub>220</sub> and G<sub>225</sub> in subdomain IX, and the amino acid motifs of subdomains  
 VIB, VIII and IX (Hardie G and Hanks S (1995) Academic Press, San Diego  
 CA).

16 The novel human Jak2 kinase, hjak2, of the present application shows  
 significant conservation of the diagnostic kinase residues which allowed  
 its identification from among the isolated cDNAs of a placenta library, the  
 anatomy and physiology of which is briefly described below.

21 The placenta is a thickened disk-shaped temporary organ that  
 interchanges gases, nutrients, hormones, excretory products, humoral  
 antibodies (IgG), and any other circulating substances between the maternal  
 and fetal bloodstreams. Receptors facilitate the transport of glucose,  
 amino acids, and IgG directly from maternal blood to fetal blood. The  
 placenta is the only organ composed of cells derived from two individuals,  
 the fetal extraembryonic chorion and the maternal endometrium. The  
 boundary between these two tissues is marked by extracellular products of  
 26 necrosis referred to as fibrinoid. This boundary results from the various  
 tissue interactions, immunological responses, etc. which occur in the  
 placenta.

31 The major tissue interaction involves the expression of paternal  
 antigens by the chorionic villi which is directly adjacent to maternal  
 blood. Although the mother initiates an immunological response, fetal  
 tissue is not typically rejected. This is attributed to the fact that the  
 fetus only expresses major histocompatibility complex (MHC) I, and not MHC  
 II which is the major cause of organ allograft rejection. In addition,  
 uterine secretions during early gestation contain significant amounts of  
 36 glucose and glycoproteins which may participate in local immunosuppression.

1 Although infections by bacteria, viruses, mycoplasmas, or parasites may  
ascend from the endocervical canal or reach the placenta through maternal  
blood, they rarely cause gross pathological changes because of maternal  
immune defense.

6 Soon after implantation, fetal villi begin to control maternal  
physiology to create an optimal environment for development. This involves  
the production of chorionic gonadotropin, estrogen and progesterone,  
chorionic somatomammotropin, insulin-like growth factors, platelet derived  
growth factor, prolactin, and various cytokines. These and other factors  
such as hjak2 certainly regulate the numerous activities (respiratory,  
11 immunological, gastrointestinal, and urinary) which occur within the  
placenta and between maternal and fetal tissues.

The anatomy and physiology of human placenta is reviewed, inter alia,  
in Benirschke and Kaufmann, (1992) Pathology of the Human Placenta,  
Springer-Verlag, New York NY, pp. 542-635; Herrera Gonzalez and Dresser  
16 (1993) Dev Comp Immunol 17(1):1-18; Mitchell et al. (1993) Placenta 14:249-  
275; Naeye (1992) Disorders of the Placenta, Fetus, and Neonate: Diagnosis  
and Clinical Significance, Moseby Year Book, St. Louis MO; and Rutanen  
(1993) Ann Med 25:343-347.

#### 21 SUMMARY

The present invention relates to a novel human Jak2 kinase and to the  
a use of the protein and its nucleic <sup>acid</sup> ~~acid~~ sequence in the study,  
a diagnosis, <sup>prevention,</sup> ~~prevention~~ and treatment of diseases. Human Jak2 kinase (hjak2)  
was first identified as a partial nucleotide sequence in Incyte Clone  
26 179527 during a computer search for nucleotide sequence alignments among  
the cDNAs of a placenta library. A modified XL-PCR procedure, specially  
designed oligonucleotides, and cDNAs of the placenta library were used to  
extend Incyte Clone 179527 to full length. The assembled nucleotide  
sequence <sup>(SEQ ID NO:1) hjak2</sup> ~~(SEQ ID NO:1)~~, <sup>(SEQ ID NO:2)</sup> ~~(SEQ ID NO:2)~~, hjak2 encodes the polypeptide <sup>(SEQ ID NO:2)</sup> ~~(SEQ ID NO:2)~~,  
a <sup>(SEQ ID NO:2)</sup> ~~(SEQ ID NO:2)~~ HJAK2. Computer search and alignment of the full length amino acid -  
31 sequence showed that HJAK2 has 92% similarity to murine Jak2 kinase  
86:1603-7),  
G (MUSPTK1; GenBank GI 409584; Wilks AF (1989) Proc Nat Acad Sci ~~86:1603-7)~~  
which in turn has 96% sequence similarity with human Jak1 kinase. These  
homologies and the conserved residues, G<sub>48</sub>, K<sub>71</sub>, E<sub>192</sub>, and D<sub>220</sub> which all lie  
36 within the catalytic domain contributed to the naming and uses of hjak2.

hjak2 (SEQ ID NO:1)

1 The complete nucleic acid sequence encoding ~~hjak2, SEQ ID NO:1~~  
disclosed herein, provides the basis for the design of antisense molecules  
useful in diminishing or eliminating expression of the genomic nucleotide  
sequence. For example, hjak2, or its oligonucleotides, fragments,  
portions, or complement, may be used in diagnostic hybridization or  
6 amplification assays of biopsied tissue to detect and/or quantify  
abnormalities in gene expression associated with an immunological disorder.  
The present invention also relates, in part, to the inclusion of the  
nucleic acid sequence in an expression vector which can be used to  
transform host cells or organisms. Such transgenic hosts are useful for  
11 production and recovery of the encoded HJAK2.

The invention further comprises using purified HJAK2 polypeptide to  
produce antibodies or to identify antagonists or inhibitors which bind  
HJAK2. Anti-HJAK antibodies may be used in membrane, tissue-based or ELISA  
technologies to detect any disease state or condition related to the  
16 aberrant expression of HJAK2. Antibodies, antagonists or inhibitors can be  
used to bind HJAK2 preventing the transfer of high energy phosphate  
molecules and therefore signal transduction. The invention also comprises  
pharmaceutical compositions containing the peptide, antibodies, antagonists  
or inhibitors for the diagnosis, prevention or treatment of conditions  
21 associated with altered or uncontrolled hjak2 expression. These conditions  
may include, but are not limited to: arteriosclerosis, asthma, bronchitis,  
emphysema, inflammatory bowel disease, leukemia, oncogenesis,  
osteoarthritis, psoriasis, rheumatoid arthritis, septic shock, and systemic  
lupus erythematosus. Steps for testing a biological sample with probes,  
26 oligomers, fragments or portions of the hjak2 nucleotide sequence or  
antibodies produced against the purified HJAK2 protein are provided.

Antisense molecules, antibodies, antagonists or inhibitors (including  
proteins, peptides, oligopeptides or organic molecules capable of  
compromising or modulating HJAK2 expression) may also be used for  
31 therapeutic purposes, for example, in neutralizing the aberrant activity  
of a HJAK2 associated with, for example, inflammation or oncogenesis. The  
present invention also provides for pharmaceutical compositions for the  
treatment of disease states associated with aberrant expression of hjak2  
comprising the forementioned antisense molecules, antibodies, antagonists  
36 or inhibitors.

## DESCRIPTION OF THE FIGURES

Figure 1 displays an alignment of the ~~nucleic acid and amino acid~~ <sup>these and in the</sup> sequences of human jak2 kinase. Alignments shown in ~~this and the~~ following figures were produced using the multisequence alignment program of DNASTAR software (DNASTAR Inc., Madison WI). Figures 2A, 2B, 2C, 2D, and 2E show ~~Figure 2 shows~~ the amino acid sequence similarity between ~~MUSPK1, GI-~~ <sup>and HAK2.</sup>

## DETAILED DESCRIPTION OF THE INVENTION

Definitions

As used herein, the abbreviation for the novel human Jak2 kinase in lower case (hjak2) refers to a gene, cDNA, ~~RNA~~ <sup>RNA</sup>, or nucleic acid ~~sequence~~ <sup>sequence</sup>, while the upper case version (HJAK2) refers to a protein, polypeptide, peptide, oligopeptide, or amino acid sequence.

An "oligonucleotide" or "oligomer" is a stretch of nucleotide residues which has a sufficient number of bases to be used in a polymerase chain reaction (PCR). These short sequences are based on (or designed from) genomic or cDNA sequences and are used to amplify, confirm, or reveal the presence of an identical, similar or complementary DNA or RNA in a particular cell or tissue. Oligonucleotides or oligomers comprise portions of a DNA sequence having at least about 10 nucleotides and as many as about 50 nucleotides, preferably about 15 to 30 nucleotides. They are chemically synthesized and may be used as probes.

"Probes" are nucleic acid sequences of variable length, preferably between at least about 10 and as many as about 6,000 nucleotides. They are used in the detection of identical, similar, or complementary nucleic acid sequences. Longer length probes are usually obtained from a natural or recombinant source, are highly specific and much slower to hybridize than oligonucleotides. They may be single- or double-stranded and are carefully designed to have specificity in PCR, hybridization membrane-based, or ELISA-like technologies.

"Reporter" molecules are chemical moieties used for labelling a nucleic or amino acid sequence. They include, but are not limited to, radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents. Reporter molecules associate with, establish the presence of, and may allow quantification of a particular nucleic or amino acid sequence.

1

A "portion" or "fragment" of a polynucleotide or nucleic acid comprises all or any part of the nucleotide sequence having fewer nucleotides than about 6 kb, preferably fewer than about 1 kb which can be used as a probe. Such probes may be labelled with reporter molecules using nick translation, Klenow fill-in reaction, PCR or other methods well known in the art. After pretesting to optimize reaction conditions and to eliminate false positives, nucleic acid probes may be used in Southern, northern or in situ hybridizations to determine whether DNA or RNA encoding the protein is present in a biological sample, cell type, tissue, organ or organism.

11

"Recombinant nucleotide variants" are polynucleotides which encode a protein. They may be synthesized by making use of the "redundancy" in the genetic code. Various codon substitutions, such as the silent changes which produce specific restriction sites or codon usage-specific mutations, may be introduced to optimize cloning into a plasmid or viral vector or expression in a particular prokaryotic or eukaryotic host system, respectively.

16

21

"Linkers" are synthesized palindromic nucleotide sequences which create internal restriction endonuclease sites for ease of cloning the genetic material of choice into various vectors. "Polylinkers" are engineered to include multiple restriction enzyme sites and provide for the use of both those enzymes which leave 5' and 3' overhangs such as BamHI, EcoRI, PstI, KpnI and Hind III or which provide a blunt end such as EcoRV, SnaBI and StuI.

26

"Control elements" or "regulatory sequences" are those nontranslated regions of the gene or DNA such as enhancers, promoters, introns and 3' untranslated regions which interact with cellular proteins to carry out replication, transcription, and translation. They may occur as boundary sequences or even split the gene. They function at the molecular level and along with regulatory genes are very important in development, growth, differentiation and aging processes.

31

"Chimeric" molecules are polynucleotides or polypeptides which are created by combining one or more of nucleotide sequences of this invention (or their parts) with additional nucleic acid sequence(s). Such combined sequences may be introduced into an appropriate vector and expressed to

36

1 give rise to a chimeric polypeptide which may be expected to be different from the native molecule in one or more of the following characteristics: cellular location, distribution, ligand-binding affinities, interchain affinities, degradation/turnover rate, signalling, etc.

6 "Active" refers to those forms, fragments, or domains of an amino acid sequence which display the biologic and/or immunogenic activity characteristic of the naturally occurring peptide.

"Naturally occurring HJAK2" refers to a polypeptide produced by cells which have not been genetically engineered or which have been genetically engineered to produce the same sequence as that naturally produced.

11 Specifically contemplated are various polypeptides which arise from post-  
~~translational~~ <sup>translational</sup> modifications. Such modifications of the polypeptide include but are not limited to acetylation, carboxylation, glycosylation, phosphorylation, lipidation and acylation.

16 "Derivative" refers to those polypeptides which have been chemically modified by such techniques as ubiquitination, labelling (see above), pegylation (derivatization with polyethylene glycol), and chemical insertion or substitution of amino acids such as ornithine which do not normally occur in human proteins.

21 "Recombinant polypeptide variant" refers to any polypeptide which differs from naturally occurring HJAK2 by amino acid insertions, deletions and/or substitutions, created using recombinant DNA techniques. Guidance in determining which amino acid residues may be replaced, added or deleted without abolishing characteristics of interest may be found by comparing the sequence of HJAK2 with that of related polypeptides and minimizing the number of amino acid sequence changes made in highly conserved regions.

26 Amino acid "substitutions" are defined as one for one amino acid replacements. They are conservative in nature when the substituted amino acid has similar structural and/or chemical properties. Examples of conservative replacements are substitution of a leucine with an isoleucine or valine, an aspartate with a glutamate, or a threonine with a serine.

31 Amino acid "insertions" or "deletions" are changes to or within an amino acid sequence. They typically fall in the range of about 1 to 5 amino acids. The variation allowed in a particular amino acid sequence may be experimentally determined by producing the peptide synthetically or by systematically making insertions, deletions, or substitutions of

nucleotides in the hjak2 sequence using recombinant DNA techniques.

A "signal or leader sequence" is a short amino acid sequence which or can be used, when desired, to direct the polypeptide through a membrane of a cell. Such a sequence may be naturally present on the polypeptides of the present invention or provided from heterologous sources by recombinant DNA techniques.

An "oligopeptide" is a short stretch of amino acid residues and may be expressed from an oligonucleotide. It may be functionally equivalent to and either the same length as or considerably shorter than <sup>a</sup>~~a "fragment", "portion", or "segment" of~~ a polypeptide. Such sequences comprise a stretch of amino acid residues of at least about 5 amino acids and often about 17 or more amino acids, typically at least about 9 to 13 amino acids, and of sufficient length to display biologic and/or immunogenic activity.

An "inhibitor" is a substance which retards or prevents a chemical or physiological reaction or response. Common inhibitors include but are not limited to antisense molecules, antibodies, antagonists and their derivatives.

A "standard" is a quantitative or qualitative measurement use for comparison. Preferably, it is based on a statistically appropriate number of samples and is created to use as a basis of comparison when performing diagnostic assays, running clinical trials, or following patient treatment profiles. The samples of a particular standard may be normal or similarly abnormal.

"Animal" as used herein may be defined to include human, domestic <sup>etc.</sup> (cats, dogs, ~~etc.~~), agricultural (cows, horses, sheep, goats, chicken, fish, <sup>etc.</sup>), ~~etc.~~ or test species (frogs, mice, rats, rabbits, <sup>simians, etc.</sup> ~~simians, etc.~~).

"Conditions" includes cancers, disorders or diseases in which hjak2 activity may be implicated. These specifically include, but are not limited to, anemia, arteriosclerosis, asthma, bronchitis, emphysema, gingivitis, inflammatory bowel disease, insulin-dependent diabetes mellitus, leukemia, multiple endocrine neoplasias, osteoarthritis, osteoporosis, pulmonary fibrosis, rheumatoid arthritis, septic shock syndromes, and systemic lupus erythematosus.

Since the list of technical and scientific terms cannot be all encompassing, any undefined terms shall be construed to have the same meaning as is commonly understood by one of skill in the art to which this



invention belongs. Furthermore, the singular forms "a", "an" and "the" include plural referents unless the context clearly dictates otherwise. For example, reference to a "restriction enzyme" or a "high fidelity enzyme" may include mixtures of such enzymes and any other enzymes fitting the stated criteria, or reference to the method includes reference to one or more methods for obtaining cDNA sequences which will be known to those skilled in the art or will become known to them upon reading this specification.

Before the present sequences, variants, formulations and methods for making and using the invention are described, it is to be understood that the invention is not to be limited only to the particular sequences, variants, formulations or methods described. The sequences, variants, formulations and methodologies may vary, and the terminology used herein is for the purpose of describing particular embodiments. The terminology and definitions are not intended to be limiting since the scope of protection will ultimately depend upon the claims.

#### Description of the Invention

The present invention provides for a purified polynucleotide which encodes a novel human Jak2 kinase which is expressed in human cells or tissues. The human Jak2 kinase (hjak2; Incyte Clone 179527) was first identified among the cDNAs from <sup>a</sup> placenta cDNA library. The naming and proscribed uses of the present invention are based in part on the conserved residues found in HJAK2. These particularly include the residues G<sub>48</sub>, K<sub>73</sub>, E<sub>192</sub>, and D<sub>220</sub>, which are all found within the catalytic domain. Computer search and alignment of the full length amino acid sequences showed that HJAK2 has 92% similarity to murine Jak2 kinase (MUSPTK1; GenBank GI 409584; Wilks AF (1989) Proc Nat Acad Sci <sup>86:1603-7</sup> ~~86:1603-7~~) which in turn has 96% sequence similarity with human Jak1 kinase.

Purified nucleotide sequences, such as hjak2, have numerous applications in techniques known to those skilled in the art of molecular biology. These techniques include their use as PCR or hybridization probes, for chromosome and gene mapping, in the production of sense or antisense nucleic acids, in screening for new therapeutic molecules, etc. These examples are well known and are not intended to be limiting. Furthermore, the nucleotide sequences disclosed herein may be used in molecular biology techniques that have not yet been developed, provided the

new techniques rely on properties of nucleotide sequences that are currently known, ~~including but not limited to~~ such properties as the triplet genetic code and specific base pair interactions.

As a result of the degeneracy of the genetic code, a multitude of HJAK2-encoding nucleotide sequences may be <sup>produced. Some</sup> ~~produced and some~~ of these will bear only minimal homology to the endogenous sequence of any known and naturally occurring Jak2 kinase sequence. This invention has specifically contemplated each and every possible variation of nucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the nucleotide sequence of naturally occurring HJAK2 and all such variations are to be considered as being specifically disclosed.

Although the hjak2 nucleotide sequence and its derivatives or variants are preferably capable of identifying the nucleotide sequence of the naturally occurring HJAK2 under optimized conditions, it may be advantageous to produce HJAK2-encoding nucleotide sequences possessing a substantially different codon usage. Codons can be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic expression host in accordance with the frequency with which particular codons are utilized by the host ~~(see)~~. Other reasons for substantially altering the nucleotide sequence encoding the HJAK2 without altering the encoded amino acid sequence include the production of RNA transcripts having more desirable properties, such as a longer half-life, than transcripts produced from the naturally occurring sequence.

Nucleotide sequences encoding HJAK2 may be joined to a variety of other nucleotide sequences by means of well established recombinant DNA techniques (Sambrook J et al (1989) Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, Chapters 4, 8, 16, and 17 (1989) Current Protocols in Molecular Biology, John Wiley & Sons, New York New York, NY, Chapters 9, 13, and 16).

~~Useful~~ Useful sequences for joining to hjak2 include an assortment of cloning vectors such as plasmids, cosmids, lambda phage derivatives, phagemids, and the like. Vectors of interest include vectors for replication, expression, probe generation, sequencing, and the like. In general, vectors of interest may contain an origin of replication functional in at least one organism, convenient restriction endonuclease sensitive sites, and selectable markers for one or more host cell systems.

1 PCR as described in US Patent Nos. 4,683,195; 4,800,195; and  
 4,965,188 provides additional uses for oligonucleotides based upon the  
 hjak2 nucleotide sequence. Such oligomers are generally chemically  
 synthesized, but they may be of recombinant origin or a mixture of both.  
 Oligomers generally comprise two nucleotide sequences, one with sense  
 6 orientation (5'→3') and one with antisense (3' to 5') employed under  
 optimized conditions for identification of a specific gene or diagnostic  
 use. The same two oligomers, nested sets of oligomers, or even a  
 degenerate pool of oligomers may be employed under less stringent  
 conditions for identification and/or quantitation of closely related DNA or  
 11 RNA sequences.

Full length genes may be cloned utilizing partial nucleotide sequence  
 and various methods known in the art. Gobinda et al (1993; PCR Methods  
 Applic 2:318-22) disclose "restriction-site PCR" as a direct method which  
 uses universal primers to retrieve unknown sequence adjacent to a known  
 16 locus. First, genomic DNA is amplified in the presence of primer to linker  
 and a primer specific to the known region. The amplified sequences are  
 subjected to a second round of PCR with the same linker primer and another  
 specific primer internal to the first one. Products of each round of PCR  
 are transcribed with an appropriate RNA polymerase and sequenced using  
 21 reverse transcriptase. Gobinda et al present data concerning Factor IX for  
 which they identified a conserved stretch of 20 nucleotides in the 3'  
 noncoding region of the gene.

Inverse PCR is the first method to report successful acquisition of  
 unknown sequences starting with primers based on a known region (Triglia T  
 26 et al (1988) Nucleic Acids Res 16:8186). The method uses several restriction  
 enzymes to generate a suitable fragment in the known region of a gene. The  
 fragment is then circularized by intramolecular ligation and used as a PCR  
 template. Divergent primers are designed from the known region. The  
 multiple rounds of restriction enzyme digestions and ligations that are  
 31 necessary prior to PCR make the procedure slow and expensive (Gobinda et  
 al, supra).

Capture PCR (Lagerstrom M et al (1991) PCR Methods Applic 1:111-19)  
 is a method for PCR amplification of DNA fragments adjacent to a known  
 sequence in human and YAC DNA. As noted by Gobinda et al (supra), capture  
 36 PCR also requires multiple restriction enzyme digestions and ligations to

place an engineered double-stranded sequence into an unknown portion of the DNA molecule before PCR. Although the restriction and ligation reactions are carried out simultaneously, the requirements for extension, immobilization and two rounds of PCR and purification prior to sequencing render the method cumbersome and time consuming.

Parker JD et al (1991; Nucleic Acids Res 19:3055-60), teach walking PCR, a method for targeted gene walking which permits retrieval of unknown sequence. In this same vein, PromoterFinder™ a new kit available from Clontech (Palo Alto CA) uses PCR and primers derived from p53 to walk in genomic DNA. Nested primers and special PromoterFinder libraries are used to detect upstream sequences such as promoters and regulatory elements. This process avoids the need to screen libraries and is useful in finding intron/exon junctions.

Another new PCR method, "Improved Method for Obtaining Full Length cDNA Sequences" by Guegler et al, Patent Application Serial No 08/487,112, filed June 7, 1995 and hereby incorporated by reference, employs XL-PCR™ (Perkin-Elmer, Foster City CA) to amplify and extend partial nucleotide sequence into longer pieces of DNA. This method was developed to allow a single researcher to process multiple genes (up to 20 or more) at one time and to obtain an extended (possibly full-length) sequence within 6-10 days. This new method replaces methods which use labelled probes to screen plasmid libraries and allow one researcher to process only about 3-5 genes in 14-40 days.

In the first step, which can be performed in about two days, any two of a plurality of primers are designed and synthesized based on a known partial sequence. In step 2, which takes about six to eight hours, the sequence is extended by PCR amplification of a selected library. Steps 3 and 4, which take about one day, are purification of the amplified cDNA and its ligation into an appropriate vector. Step 5, which takes about one day, involves transforming and growing up host bacteria. In step 6, which takes approximately five hours, PCR is used to screen bacterial clones for extended sequence. The final steps, which take about one day, involve the preparation and sequencing of selected clones.

If the full length cDNA has not been obtained, the entire procedure is repeated using either the original library or some other preferred library. The preferred library may be one that has been size-selected to

1 include only larger cDNAs or may consist of single or combined commercially  
available libraries, eg. lung, liver, heart and brain from Gibco/BRL  
(Gaithersburg MD). The cDNA library may have been prepared with oligo (dT)  
or random priming. Random primed libraries are preferred in that they will  
6 contain more sequences which contain 5' ends of genes. A randomly primed  
library may be particularly useful if an oligo (dT) library does not yield  
a complete gene. It must be noted that the larger and more complex the  
protein, the less likely it is that the complete gene will be found in a  
single plasmid.

11 A new method for analyzing either the size or the nucleotide sequence  
of PCR products is capillary electrophoresis. Systems for rapid sequencing  
are available from Perkin Elmer (Foster City CA), Beckman Instruments  
(Fullerton CA), and other companies. Capillary sequencing employs flowable  
polymers for electrophoretic separation, four different fluorescent dyes  
(one for each nucleotide) which are laser activated, and detection of the  
16 emitted wavelengths by a charge coupled device camera. Output/light  
intensity is converted to electrical signal using appropriate software (eg.  
Genotyper™ and Sequence Navigator™ from Perkin Elmer) and the entire  
process from loading of samples to computer analysis and electronic data  
display is computer controlled. Capillary electrophoresis provides greater  
21 resolution and is many times faster than standard gel based procedures. It  
is particularly suited to the sequencing of small pieces of DNA which might  
be present in limited amounts in a particular sample. The reproducible  
sequencing of up to 350 bp of M13 phage DNA in 30 min has been reported  
(Ruiz-Martinez MC et al (1993) Anal Chem 65:2851-8).

26 Another aspect of the subject invention is to provide for hjak2  
hybridization probes which are capable of hybridizing with naturally  
occurring nucleotide sequences encoding HJAK2. The stringency of the  
hybridization conditions will determine whether the probe identifies only  
the native nucleotide sequence of hjak2 or sequences of other closely  
31 related Jak2 kinase molecules. If degenerate hjak2 nucleotide sequences of  
the subject invention are used for the detection of related kinase encoding  
sequences, they should preferably contain at least 50% of the nucleotides  
of the sequences presented herein. Hybridization probes of the subject  
invention may be derived from the nucleotide sequence presented in SEQ ID  
36 NO 1 or from surrounding genomic sequences comprising untranslated regions

1 such as promoters, enhancers and introns. Such hybridization probes may be labelled with appropriate reporter molecules.

Means for producing specific hybridization probes for this Jak2 kinase include oligolabelling, nick translation, end-labelling or PCR amplification using a labelled nucleotide. Alternatively, the cDNA  
6 sequence may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3 or SP6 and labelled nucleotides. A number of companies (such as Pharmacia Biotech, Piscataway NJ; Promega, Madison WI; US Biochemical Corp, Cleveland, OH; etc.) supply commercial kits and  
11 protocols for these procedures.

It is also possible to produce a DNA sequence, or portions thereof, entirely by synthetic chemistry. Sometimes the source of information for producing this sequence comes from the known homologous sequence from  
16 closely related organisms. After synthesis, the nucleic acid sequence can be used alone or joined with a pre-existing sequence and inserted into one of the many available DNA vectors and their respective host cells using techniques well known in the art. Moreover, synthetic chemistry may be used to introduce specific mutations into the nucleotide sequence.  
21 Alternatively, a portion of sequence in which a mutation is desired can be synthesized and recombined with a portion of an existing genomic or recombinant sequence.

Hjak2 nucleotide sequence can be used in a diagnostic test or assay to detect disorder or disease processes associated with abnormal expression  
26 of hjak2. The nucleotide sequence is added to a sample (fluid, cell or tissue) from a patient under hybridizing conditions. After an incubation period, the sample is washed with a compatible fluid which optionally contains a reporter molecule which will bind the specific nucleotide. After the compatible fluid is rinsed off, the reporter molecule is  
31 quantitated and compared with a standard for that fluid, cell or tissue. If hjak2 expression is significantly different from the standard, the assay indicates the presence of disorder or disease. The form of such qualitative or quantitative methods may include northern analysis, dot blot or other membrane-based technologies, dip stick, pin or chip technologies,  
36 PCR, ELISAs or other multiple sample format technologies.

1 This same assay, combining a sample with the nucleotide sequence, is  
applicable in evaluating the efficacy of a particular therapeutic treatment  
regime. It may be used in animal studies, in clinical trials, or in  
monitoring the treatment of an individual patient. First, standard  
expression must be established for use as a basis of comparison. Second,  
6 samples from the animals or patients affected by a disorder or disease are  
combined with the nucleotide sequence to evaluate the deviation from the  
standard or normal profile. Third, an entirely new or pre-existing  
therapeutic agent is administered, and a treatment profile is generated.  
This post-treatment assay is evaluated to determine whether the patient  
11 profile progresses toward or returns to the standard pattern. Successive  
treatment profiles may be used to show the efficacy of treatment over a  
period of several days or several months.

The nucleotide sequence for *hjak2* can also be used to generate probes  
for mapping native genomic sequence. The sequence may be mapped to a  
16 particular chromosome or to a specific region of the chromosome using well  
known techniques. These include in situ hybridization to chromosomal  
spreads (Verma et al (1988) Human Chromosomes: A Manual of Basic  
Techniques, Pergamon Press, New York City), flow-sorted chromosomal  
preparations, or artificial chromosome constructions such as yeast  
21 artificial chromosomes (YACs), bacterial artificial chromosomes (BACs),  
bacterial P1 constructions or single chromosome cDNA libraries.

In situ hybridization of chromosomal preparations and physical  
mapping techniques such as linkage analysis using established chromosomal  
markers are invaluable in extending genetic maps. Examples of such genetic  
26 maps can regularly be found in the journal Science (eg, 1994; 265:1981f).  
Often the placement of a gene on the chromosome of another mammalian  
species may reveal associated markers even if the number or arm of a  
particular human chromosome is not known. New sequences can be assigned to  
chromosomal arms, or parts thereof, by physical mapping. This provides  
31 valuable information to investigators searching for disease genes using  
positional cloning or other gene discovery techniques. Once a disease or  
syndrome, such as ataxia telangiectasia (AT), has been crudely localized by  
genetic linkage to a particular genomic region, for example, AT to 11q22-23  
(Gatti et al (1988) Nature 336:577-580), any sequences mapping to that area  
36 may represent associated or regulatory genes for further investigation.

1 The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc. between normal and carrier or affected individuals.

6 The nucleotide sequence encoding HJAK2 may be used to produce an amino acid sequence using well known methods of recombinant DNA technology. Goeddel (1990, Gene Expression Technology, Methods and Enzymology, Vol 185, Academic Press, San Diego CA) is one among many publications which teach expression of an isolated, purified nucleotide sequence. The amino acid or peptide may be expressed in a variety of host cells, either prokaryotic or eukaryotic. Host cells may be from the same species from which the nucleotide sequence was derived or from a different species. Advantages of producing an amino acid sequence or peptide by recombinant DNA technology include obtaining adequate amounts for purification and the availability of simplified purification procedures.

16 Cells transformed with hjak2 nucleotide sequence may be cultured under conditions suitable for the expression and recovery of peptide from cell culture. The peptide produced by a recombinant cell may be secreted or may be contained intracellularly depending on the sequence and/or the vector used. In general, it is more convenient to prepare recombinant proteins in secreted form, and this is accomplished by ligating hjak2 to a recombinant nucleotide sequence which directs its movement through a particular prokaryotic or eukaryotic cell membrane. Other recombinant constructions may join hjak2 to nucleotide sequence encoding a polypeptide domain which will facilitate protein purification (Kroll DJ et al (1993) DNA Cell Biol 12:441-53).

26 Direct peptide synthesis using solid-phase techniques (Creighton (1983) Proteins Structures And Molecular Principles, WH Freeman and Co, New York NY pp. 50-60) is an alternative to recombinant or chimeric peptide production. Automated synthesis may be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer in accordance with the instructions provided by the manufacturer. Additionally HJAK2 or any part thereof may be mutated during direct synthesis and combined using chemical methods with other kinase sequences, or parts thereof.

36 Although an amino acid sequence or oligopeptide used for antibody induction does not require biological activity, it must be immunogenic. HJAK2 used to induce specific antibodies may have an amino acid sequence



1 consisting of at least five amino acids and preferably at least 10 amino  
acids. Short stretches of amino acid sequence may be fused with those of  
another protein such as keyhole limpet hemocyanin, and the chimeric peptide  
used for antibody production. Alternatively, the peptide may be of  
sufficient length to contain an entire domain.

6 Antibodies specific for HJAK2 may be produced by inoculation of an  
appropriate animal with an antigenic fragment of the peptide. An antibody  
is specific for HJAK2 if it is produced against an epitope of the  
polypeptide and binds to at least part of the natural or recombinant  
protein. Antibody production includes not only the stimulation of an  
11 immune response by injection into animals, but also analogous processes  
such as the production of synthetic antibodies, the screening of  
recombinant immunoglobulin libraries for specific-binding molecules  
(Orlandi R et al (1989) PNAS 86:3833-3837, or Huse WD et al (1989) Science  
256:1275-1281), or the in vitro stimulation of lymphocyte populations.  
16 Current technology (Winter G and Milstein C (1991) Nature 349:293-299)  
provides for a number of highly specific binding reagents based on the  
principles of antibody formation. These techniques may be adapted to  
produce molecules which specifically bind HJAK2. Antibodies or other  
appropriate molecules generated against a specific immunogenic peptide  
21 fragment or oligopeptide can be used in Western analysis, enzyme-linked  
immunosorbent assays (ELISA) or similar tests to establish the presence of  
or to quantitate amounts of HJAK2 active in normal, diseased, or  
therapeutically treated cells or tissues.

26 The examples below are provided to illustrate the subject invention.  
These examples are provided by way of illustration and are not included for  
the purpose of limiting the invention.

#### EXAMPLES

##### 31 I Placenta cDNA Library Construction

Library was constructed from normal placenta obtained from the Mayo  
Clinic. The tissue was lysed in a buffer containing guanidinium  
isothiocyanate. The lysate was extracted with phenol chloroform and  
precipitated with ethanol. Poly A<sup>+</sup> RNA was isolated using biotinylated  
36 oligo d(T) primer and streptavidin coupled to a paramagnetic particle

(Promega Corp. Madison WI) and sent to Stratagene (La Jolla CA) for cDNA library preparation. The cDNA synthesis was primed using both oligo d(T) and random hexamers, and the two cDNA libraries were treated separately. Synthetic adapter oligonucleotides were ligated onto the ends of the cDNAs which were digested with XhoI and inserted into the Uni-ZAP™ vector system (Stratagene).

The pBluescript™ phagemid (Stratagene) was excised from each library, and phagemids from the two cDNA libraries were combined into a single library by mixing equal numbers of bacteriophage. The phagemids were transformed into *E. coli* host strain XL1-Blue™ (Stratagene). Enzymes from both pBluescript and a cotransformed f1 helper phage nicked the DNA, initiated new DNA synthesis, and created the smaller, single-stranded circular plasmid DNA molecules which contained the cDNA insert. The plasmid DNA was released, purified, and used to reinfect fresh host cells (SOLR, Stratagene). Presence of the  $\beta$ -lactamase gene on the plasmid allowed transformed bacteria to grow on medium containing ampicillin.

## II Isolation of cDNA Clones

Plasmid DNAs containing the cDNA insert were purified using the QIAWELL-8 Plasmid Purification System from QIAGEN Inc (Chatsworth CA) according to standard protocol. The DNA was eluted and prepared for DNA sequencing and other analytical manipulations.

The cDNA inserts from random isolates of the placenta library were partially sequenced. The cDNAs were sequenced by the method of Sanger F and AR Coulson (1975; J Mol Biol 94:441f), using a Catalyst 800 or a Hamilton Micro Lab 2200 (Hamilton, Reno NV) in combination with four Peltier Thermal Cyclers (PTC200 from MJ Research, Watertown MA) and Applied Biosystems 377 or 373 DNA Sequencing Systems (Perkin Elmer), and reading frame was determined.

## III Sequencing of cDNA Clones

The cDNA inserts from random isolates of the placenta library were sequenced in part. Methods for DNA sequencing are well known in the art and employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE® (US Biochemical Corp) or Taq polymerase. Methods to extend the DNA from an oligonucleotide primer annealed to the DNA template of interest have been developed for both single- and double-stranded templates. Chain

1 termination reaction products were separated using electrophoresis and  
detected via their incorporated, labelled precursors. Recent improvements  
in mechanized reaction preparation, sequencing and analysis have permitted  
expansion in the number of sequences that can be determined per day.  
Preferably, the process is automated with machines such as the Hamilton  
6 Micro Lab 2200 (Hamilton, Reno NV), Peltier Thermal Cycler (PTC200; MJ  
Research, Watertown MA) and the Applied Biosystems Catalyst 800 and 377 and  
373 DNA sequencers.

11 The quality of any particular cDNA library may be determined by  
performing a pilot scale analysis of the cDNAs and checking for percentages  
of clones containing vector, lambda or *E. coli* DNA, mitochondrial or  
repetitive DNA, and clones with exact or homologous matches to public  
databases. The number of unique sequences, those having no known match in  
any available database, are then recorded.

#### 16 IV Homology Searching of cDNA Clones and Their Deduced Proteins

Each sequence so obtained was compared to sequences in GenBank using  
a search algorithm developed by Applied Biosystems and incorporated into  
the INHERIT<sup>®</sup> 670 Sequence Analysis System. In this algorithm, Pattern  
Specification Language (TRW Inc, Los Angeles CA) was used to determine  
21 regions of homology. The three parameters that determine how the sequence  
comparisons run were window size, window offset, and error tolerance.  
Using a combination of these three parameters, the DNA database was  
searched for sequences containing regions of homology to the query  
sequence, and the appropriate sequences were scored with an initial value.  
26 Subsequently, these homologous regions were examined using dot matrix  
homology plots to distinguish regions of homology from chance matches.  
Smith-Waterman alignments were used to display the results of the homology  
search.

31 Peptide and protein sequence homologies were ascertained using the  
INHERIT<sup>®</sup> 670 Sequence Analysis System in a way similar to that used in DNA  
sequence homologies. Pattern Specification Language and parameter windows  
were used to search protein databases for sequences containing regions of  
homology which were scored with an initial value. Dot-matrix homology  
plots were examined to distinguish regions of significant homology from  
36 chance matches.

Alternatively, BLAST, which stands for Basic Local Alignment Search

1 Tool, is used to search for local sequence alignments (Altschul SF (1993) J  
Mol Evol 36:290-300; Altschul, SF et al (1990) J Mol Biol 215:403-10).  
BLAST produces alignments of both nucleotide and amino acid sequences to  
determine sequence similarity. Because of the local nature of the  
alignments, BLAST is especially useful in determining exact matches or in  
6 identifying homologs. While it is useful for matches which do not contain  
gaps, it is inappropriate for performing motif-style searching. The  
fundamental unit of BLAST algorithm output is the High-scoring Segment Pair  
(HSP).

11 An HSP consists of two sequence fragments of arbitrary but equal  
lengths whose alignment is locally maximal and for which the alignment  
score meets or exceeds a threshold or cutoff score set by the user. The  
BLAST approach is to look for HSPs between a query sequence and a database  
sequence, to evaluate the statistical significance of any matches found,  
and to report only those matches which satisfy the user-selected threshold  
of significance. The parameter E establishes the statistically significant  
16 threshold for reporting database sequence matches. E is interpreted as  
the upper bound of the expected frequency of chance occurrence of an HSP  
(or set of HSPs) within the context of the entire database search. Any  
database sequence whose match satisfies E is reported in the program  
21 output.

The partial hjak2 molecule presented and claimed in this application  
was identified using the criteria above. The full length nucleic and amino  
acid sequences for this novel human Jak2 kinase are shown in Fig 1. Fig 2  
shows the alignment between the translated amino acid sequence for hjak2  
26 and the closest related molecule, murine Jak2 kinase (MUSPTK1; GenBank GI  
409584; Wilks AF (1989) Proc Nat Acad Sci 86:1603-7).

#### V Extension of cDNAs to Full Length

31 The partial sequence originally identified in Incyte Clone 179527 was  
used to design oligonucleotide primers for extension of the cDNAs to full  
length. Primers are designed based on known sequence; one primer is  
synthesized to initiate extension in the antisense direction (XLR) and the  
other to extend sequence in the sense direction (XLF). The primers allow  
the sequence to be extended "outward" generating amplicons containing new,  
36 unknown nucleotide sequence for the gene of interest. The primers may be  
designed using Oligo 4.0 (National Biosciences Inc, Plymouth MN), or

another appropriate program, to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68°-72° C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

The placenta cDNA library was used with XLR = GGGCGGAAGTGCTCTCGGCGGAAG and XLF = AGTGTGCTACAGTGCTGGTCGTCG primers to extend and amplify Incyte Clone 179527 to obtain the full length Jak2 kinase sequence.

By following the instructions for the XL-PCR kit and thoroughly mixing the enzyme and reaction mix, high fidelity amplification is obtained. Beginning with 40 pmol of each primer and the recommended concentrations of all other components of the kit, PCR is performed using the Peltier Thermal Cycler (PTC200; MJ Research, Watertown MA) and the following parameters:

Step 1	94° C for 1 min (initial denaturation)
Step 2	65° C for 1 min
Step 3	68° C for 6 min
Step 4	94° C for 15 sec
Step 5	65° C for 1 min
Step 6	68° C for 7 min
Step 7	Repeat step 4-6 for 15 additional cycles
Step 8	94° C for 15 sec
Step 9	65° C for 1 min
Step 10	68° C for 7:15 min
Step 11	Repeat step 8-10 for 12 cycles
Step 12	72° C for 8 min
Step 13	4° C (and holding)

A 5-10 µl aliquot of the reaction mixture is analyzed by electrophoresis on a low concentration (about 0.6-0.8%) agarose mini-gel to determine which reactions were successful in extending the sequence. Although all extensions potentially contain a full length gene, some of the largest products or bands are selected and cut out of the gel. Further purification involves using a commercial gel extraction method such as QIAQuick™ (QIAGEN Inc). After recovery of the DNA, Klenow enzyme is used to trim single-stranded, nucleotide overhangs creating blunt ends which facilitate religation and cloning.

After ethanol precipitation, the products are redissolved in 13 µl of ligation buffer. Then, 1µl T4-DNA ligase (15 units) and 1µl T4 polynucleotide kinase are added, and the mixture is incubated at room temperature for 2-3 hours or overnight at 16° C. Competent E. coli cells

(in 40  $\mu$ l of appropriate media) are transformed with 3  $\mu$ l of ligation mixture and cultured in 80  $\mu$ l of SOC medium (Sambrook J et al, supra). After incubation for one hour at 37° C, the whole transformation mixture is plated on Luria Bertani (LB)-agar (Sambrook J et al, supra) containing 2xCarb. The following day, 12 colonies are randomly picked from each plate and cultured in 150  $\mu$ l of liquid LB/2xCarb medium placed in an individual well of an appropriate, commercially-available, sterile 96-well microtiter plate. The following day, 5  $\mu$ l of each overnight culture is transferred into a non-sterile 96-well plate and after dilution 1:10 with water, 5  $\mu$ l of each sample is transferred into a PCR array.

For PCR amplification, 18  $\mu$ l of concentrated PCR reaction mix (3.3x) containing 4 units of rTth DNA polymerase, a vector primer and one or both of the gene specific primers used for the extension reaction are added to each well. Amplification is performed using the following conditions:

Step 1	94° C for 60 sec
Step 2	94° C for 20 sec
Step 3	55° C for 30 sec
Step 4	72° C for 90 sec
Step 5	Repeat steps 2-4 for an additional 29 cycles
Step 6	72° C for 180 sec
Step 7	4° C (and holding)

Aliquots of the PCR reactions are run on agarose gels together with molecular weight markers. The sizes of the PCR products are compared to the original partial cDNAs, and appropriate clones are selected, ligated into plasmid and sequenced.

## VI Diagnostic Assay Using Hjak2 Specific Oligomers

In those cases where a specific condition (see definitions, supra) is suspected to involve expression of altered quantities of hjak2, oligomers may be designed to establish the presence and/or quantity of mRNA expressed in a biological sample. There are several methods currently being used to quantitate the expression of a particular molecule. Most of these methods use radiolabelled (Melby PC et al 1993 J Immunol Methods 159:235-44) or biotinylated (Duplaa C et al 1993 Anal Biochem 229-36) nucleotides, coamplification of a control nucleic acid, and standard curves onto which the experimental results are interpolated. Quantitation may be speeded up by running the assay in an ELISA format where the oligomer-of-interest is presented in various dilutions and a colorimetric response gives rapid

1 quantitation. For example, a complete HJAK2 deficiency may result in the  
inability to undergo cell division or to react to an infectious organism.  
In like manner, overexpression may cause major inflammation, swelling and  
major tissue damage. In either case, a quick diagnosis may allow health  
professionals to treat the condition and prevent worsening of the  
6 condition. This same assay can be used to monitor progress of the patient  
as his/her physiological situation moves toward the normal range during  
therapy.

#### VII Sense or Antisense Molecules

11 Knowledge of the correct cDNA sequence of this Jak2 kinase or its  
regulatory elements enable its use as a tool in sense (Yousoufian H and HF  
Lodish 1993) Mol Cell Biol 13:98-104) or antisense (Eguchi et al (1991)  
Annu Rev Biochem 60:631-652) technologies for the investigation or  
alteration of gene expression. To inhibit in vivo or in vitro cdp  
16 expression, an oligonucleotide based on the coding sequence of an hjak2  
designed with Oligo 4.0 (National Biosciences Inc) is used. Alternatively,  
a fragment of an hjak2 is produced by digesting hjak2 coding sequence with  
restriction enzymes. These enzymes and specific restrictions sites may be  
selected using Inherit Analysis software (Applied Biosystems), and the  
21 strands separated by heating the fragments and selecting for the antisense  
strand. Either the oligonucleotide or the fragment may be used to inhibit  
hjak2 expression. Furthermore, antisense molecules can be designed to  
inhibit promoter binding in the upstream nontranslated leader or at various  
sites along the hjak2 coding region. Alternatively, antisense molecules  
26 may be designed to inhibit translation of an mRNA into polypeptide by  
preparing an oligomer or fragment which will bind in the region spanning  
approximately -10 to +10 nucleotides at the 5' end of the coding sequence.  
These technologies are now well known to those of in the art.

out  
Dy

31 In addition to using antisense molecules constructed to interrupt  
transcription of the open reading frame, modifications of gene expression  
can be obtained by designing antisense sequences to enhancers, introns, or  
even to trans-acting regulatory genes. Similarly, inhibition can be  
achieved using Hogeboom base-pairing methodology, also known as "triple  
helix" base pairing. Triple helix pairing compromises the ability of the  
36 double helix to open sufficiently for the binding of polymerases,

transcription factors, or regulatory molecules.

Any of these types of antisense molecules may be placed in expression vectors and used to transform preferred cells or tissues. This may include introduction of the expression vector into a organ, tumor, synovial cavity or the vascular system for transient or short term therapy or introduction via gene therapy technologies for long term treatment. Transient expression may last for a month or more with a non-replicating vector and three months or more if appropriate replication elements are used in the transformation or expression system.

Stable transformation of appropriate dividing cells with a vector containing the antisense molecule can produce a transgenic cell line, tissue or organism (see, for example, Trends in Biotechnol 11:155-215 (1993) and US Patent No. 4,736,866, 12 April 1988). Those cells which assimilate or replicate enough copies of the vector to allow stable integration will also produce enough antisense molecules to compromise or entirely eliminate normal activity of the hjak2. Frequently, the function of an hjak2 can be ascertained by observing behaviors such as lethality, loss of a physiological pathway, changes in morphology, etc. at the cellular, tissue or organismal level.

#### VIII Expression of HJAK2

Expression of the HJAK2 may be accomplished by subcloning the cDNA into appropriate vectors and transfecting the vectors into host cells. In this case, the cloning vector previously used for the generation of the tissue library also provides for direct expression of the hjak2 sequence in *E. coli*. Upstream of the cloning site, this vector contains a promoter for  $\beta$ -galactosidase, followed by sequence containing the amino-terminal Met and the subsequent 7 residues of  $\beta$ -galactosidase. Immediately following these eight residues is a bacteriophage promoter useful for transcription and a linker containing a number of unique restriction sites.

Induction of an isolated, transfected bacterial strain with IPTG using standard methods will produce a fusion protein corresponding to the first seven residues of  $\beta$ -galactosidase, about 5 to 15 residues which correspond to linker, and the peptide encoded within the hjak2 cDNA. Since cDNA clone inserts are generated by an essentially random process, there is one chance in three that the included cDNA will lie in the correct frame for proper translation. If the cDNA is not in the proper reading frame, it



1 can be obtained by deletion or insertion of the appropriate number of bases  
by well known methods including in vitro mutagenesis, digestion with  
exonuclease III or mung bean nuclease, or oligonucleotide linker inclusion.

6 The cDNA can be shuttled into other vectors known to be useful for  
expression of protein in specific hosts. Oligonucleotide linkers  
containing cloning sites as well as a stretch of DNA sufficient to  
hybridize to the end of the target cDNA (25 bases) can be synthesized  
chemically by standard methods. These primers can then used to amplify the  
desired gene fragments by PCR. The resulting fragments can be digested  
11 with appropriate restriction enzymes under standard conditions and isolated  
by gel electrophoresis. Alternatively, similar gene fragments can be  
produced by digestion of the cDNA with appropriate restriction enzymes and  
filling in the missing gene sequence with chemically synthesized  
oligonucleotides. Partial nucleotide sequence from more than one kinase  
16 homolog can be ligated together and cloned into appropriate vectors to  
optimize expression.

Suitable expression hosts for such chimeric molecules include but are  
not limited to mammalian cells such as Chinese Hamster Ovary (CHO) and  
human 293 cells, insect cells such as Sf9 cells, yeast cells such as  
21 Saccharomyces cerevisiae, and bacteria such as E. coli. For each of these  
cell systems, a useful expression vector may also include an origin of  
replication to allow propagation in bacteria and a selectable marker such  
as the  $\beta$ -lactamase antibiotic resistance gene to allow selection in  
bacteria. In addition, the vectors may include a second selectable marker  
26 such as the neomycin phosphotransferase gene to allow selection in  
transfected eukaryotic host cells. Vectors for use in eukaryotic  
expression hosts may require RNA processing elements such as 3'  
polyadenylation sequences if such are not part of the cDNA of interest.

31 If native promoters are not part of the cDNA, other host specific  
promoters may be specifically combined with the coding region of hjak2.  
They include MMTV, SV40, and ~~metallothionein~~ <sup>metallothionein</sup> promoters for CHO cells; trp,  
lac, tac and T7 promoters for bacterial hosts; and alpha factor, alcohol  
oxidase and PGH promoters for yeast. In addition, transcription enhancers,  
such as the rous sarcoma virus (RSV) enhancer, may be used in mammalian  
36 host cells. Once homogeneous cultures of recombinant cells are obtained

1 through standard culture methods, large quantities of recombinantly  
produced peptide can be recovered from the conditioned medium and analyzed  
using methods known in the art.

#### IX Isolation of Recombinant HJAK2

6 HJAK2 may be expressed as a recombinant protein with one or more  
additional polypeptide domains added to facilitate protein purification.  
Such purification facilitating domains include, but are not limited to,  
metal chelating peptides such as histidine- tryptophan modules that allow  
purification on immobilized metals, protein A domains that allow  
11 purification on immobilized immunoglobulin, and the domain utilized in the  
FLAGS extension/affinity purification system (Immunex Corp, Seattle WA).  
The inclusion of a cleavable linker sequence such as Factor XA or  
enterokinase (Invitrogen) between the purification domain and the hjak2  
sequence may be useful to facilitate expression of HJAK2.

#### X Testing HJAK2 Activity

16 The sequence for HJAK2 in this application present many different  
domains (and subdomains as detailed in the background of the invention)  
which may be utilized: 1) individually for the production of antibodies, 2)  
21 in functional groups (eg. to span a membrane), and 3) as interchangeable,  
usable parts of a chimeric kinase. For example, a known, full length  
kinase such as the hjak2 kinase of this application may be used to swap  
related portions of the nucleic acid sequence, analogous to domains or  
subdomains of MAP kinase polypeptides. The chimeric nucleotides, so  
26 produced, may be introduced into prokaryotic host cells (as reviewed in  
Strosberg AD and Marullo S (1992) Trends Pharma Sci 13:95-98) or eukaryotic  
host cells. These host cells are then employed in procedures to determine  
what molecules activate the kinase or what molecules are activated by a  
kinase. Such activating or activated molecules may be of extracellular,  
31 intracellular, biologic or chemical origin.

An example of a test system, in this case for hjak2 kinase, can be  
based on the interaction of protein tyrosine kinases with chemokine  
receptors (Taniguchi T (1995) Science 268:251-255). These receptors are  
capable of activating a variety of nonreceptor protein tyrosine kinases  
36 when stimulated by an extracellular chemokine. C-X-C chemokines such as

platelet factor 4, interleukin-8, connective tissue activating protein III, neutrophil activating peptide 2, are soluble activators of neutrophils.

A standard measure of neutrophil activation involves measuring the mobilization of  $Ca^{++}$  as part of the signal transduction pathway. The experiment involves several steps. First, blood cells obtained from venipuncture are fractionated by centrifugation on density gradients. Enriched populations of neutrophils are further fractionated on columns by negative selection using antibodies specific for other blood cells types. Next, neutrophils are transformed with an expression vector containing the kinase nucleic acid sequence of interest and preloaded fluorescent probe whose emission characteristics have been altered by  $Ca^{++}$  binding. Or in the alternative, the neutrophil is preloaded with the purified kinase of interest and fluorescent probe. Then, when the cells are exposed to an appropriate chemokine, the chemokine receptor activates the kinase which, in turn, initiates  $Ca^{++}$  flux.  $Ca^{++}$  mobilization is observed and measured using fluorometry as has been described in Grynkiewicz G et al (1985) J Biol Chem 260:3440, and McColl S et al (1993) J Immunol 150:4550-4555, incorporated herëin by reference.

#### XI Identification of or Production of HJAK2 Specific Antibodies

Purified HJAK2 is used to screen a pre-existing antibody library or to raise antibodies using either polyclonal or monoclonal methodology. In a polyclonal approach, denatured protein from the reverse phase HPLC separation is obtained in quantities up to 75 mg. This denatured protein can be used to immunize mice or rabbits using standard protocols; about 100 micrograms are adequate for immunization of a mouse, while up to 1 mg might be used to immunize a rabbit. For identifying mouse hybridomas, the denatured protein can be radioiodinated and used to screen potential murine B-cell hybridomas for those which produce antibody. This procedure requires only small quantities of protein, such that 20 mg would be sufficient for labeling and screening of several thousand clones.

In a monoclonal approach, the amino acid sequence of HJAK2, as deduced from translation of the cDNA, is analyzed to determine regions of high immunogenicity. Oligopeptides comprising appropriate hydrophilic regions, as shown in Fig. 3, are synthesized and used in suitable immunization protocols to raise antibodies. Analysis to select appropriate epitopes is described by Ausubel FM et al (supra). The optimal amino acid

1 sequences for immunization are usually at the C-terminus, the N-terminus  
and those intervening, hydrophilic regions of the polypeptide which are  
likely to be exposed to the external environment when the protein is in its  
natural conformation.

6 Typically, selected peptides, about 15 residues in length, are  
synthesized using an Applied Biosystems Peptide Synthesizer Model 431A  
using fmoc-chemistry and coupled to keyhole limpet hemocyanin (KLH, Sigma)  
by reaction with M-maleimidobenzoyl- N-hydroxysuccinimide ester (MBS;  
Ausubel FM et al, supra). If necessary, a cysteine may be introduced at  
the N-terminus of the peptide to permit coupling to KLH. Rabbits are  
11 immunized with the peptide-KLH complex in complete Freund's adjuvant. The  
resulting antisera are tested for antipeptide activity by binding the  
peptide to plastic, blocking with 1% BSA, reacting with antisera, washing  
and reacting with labeled (radioactive or fluorescent), affinity purified,  
specific goat anti-rabbit IgG.

16 Hybridomas may also be prepared and screened using standard  
techniques. Hybridomas of interest are detected by screening with labeled  
HJAK2 to identify those fusions producing the monoclonal antibody with the  
desired specificity. In a typical protocol, wells of plates (FAST; Becton-  
Dickinson, Palo Alto, CA) are coated with affinity purified, specific  
21 rabbit-anti-mouse antibodies (or suitable anti-species Ig) at 10 mg/ml.  
The coated wells are blocked with 1% BSA, washed and exposed to  
supernatants from hybridomas. After incubation the wells are exposed to  
labeled HJAK2, 1 mg/ml. Clones producing antibodies will bind a quantity  
of labeled HJAK2 which is detectable above background. Such clones are  
26 expanded and subjected to 2 cycles of cloning at limiting dilution (1  
cell/3 wells). Cloned hybridomas are injected into pristine mice to  
produce ascites, and monoclonal antibody is purified from mouse ascitic  
fluid by affinity chromatography on Protein A. Monoclonal antibodies with  
affinities of at least  $10^8$  /M, preferably  $10^9$  to  $10^{10}$  or stronger, will  
31 typically be made by standard procedures as described in Harlow and Lane  
(1988) Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold  
Spring Harbor NY; and in Goding (1986) Monoclonal Antibodies: Principles  
and Practice, Academic Press, New York City, both incorporated herein by  
reference.

## XII Diagnostic Test Using HJAK2 Specific Antibodies

Particular HJAK2 antibodies are useful for the diagnosis of prepathologic conditions, and chronic or acute diseases which are characterized by differences in the amount or distribution of HJAK2. To date, HJAK2 has been found only in the placenta library; however, its activity there is most probably associated with organ function, inflammation or defense.

Diagnostic tests for HJAK2 include methods utilizing the antibody and a label to detect HJAK2 in human body fluids, tissues or extracts of such tissues. The polypeptides and antibodies of the present invention may be used with or without modification. Frequently, the polypeptides and antibodies will be labeled by joining them, either covalently or noncovalently, with a reporter molecule. A wide variety of labels and conjugation techniques are known and have been reported extensively in both the scientific and patent literature. Suitable reporter molecules or labels include those radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents previously mentioned as well as substrates, cofactors, inhibitors, magnetic particles and the like. Patents teaching the use of such labels include US Patent Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241. Also, recombinant immunoglobulins may be produced as shown in US Patent No. 4,816,567, incorporated herein by reference.

A variety of protocols for measuring soluble or membrane-bound HJAK2, using either polyclonal or monoclonal antibodies specific for the respective protein are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA) and fluorescent activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on HJAK2 is preferred, but a competitive binding assay may be employed. These assays are described, among other places, in Maddox, DE et al (1983, J Exp Med 158:1211).

## XIII Purification of Native HJAK2 Using Specific Antibodies

Native or recombinant HJAK2 can be purified by immunoaffinity chromatography using antibodies specific for that particular HJAK2. In general, an immunoaffinity column is constructed by covalently coupling the

1 anti-HJAK2 antibody to an activated chromatographic resin.

Polyclonal immunoglobulins are prepared from immune sera either by precipitation with ammonium sulfate or by purification on immobilized Protein A (Pharmacia Biotech). Likewise, monoclonal antibodies are prepared from mouse ascites fluid by ammonium sulfate precipitation or chromatography on immobilized Protein A. Partially purified immunoglobulin is covalently attached to a chromatographic resin such as CnBr-activated Sepharose (Pharmacia Biotech). The antibody is coupled to the resin, the resin is blocked, and the derivative resin is washed according to the manufacturer's instructions.

11 Such immunoaffinity columns may be utilized in the purification of HJAK2 by preparing a fraction from cells containing HJAK2 in a soluble form. This preparation may be derived by solubilization of whole cells or of a subcellular fraction obtained via differential centrifugation (with or without addition of detergent) or by other methods well known in the art. Alternatively, soluble HJAK2 containing a signal sequence may be secreted in useful quantity into the medium in which the cells are grown.

A soluble HJAK2-containing preparation is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of HJAK2 (eg, high ionic strength buffers in the presence of detergent). Then, the column is eluted under conditions that disrupt antibody/HJAK2 binding (eg, a buffer of pH 2-3 or a high concentration of a chaotrope such as urea or thiocyanate ion), and HJAK2 is collected.

#### 26 XIV. Drug Screening

This invention is particularly useful for screening therapeutic compounds by using binding fragments of HJAK2 in any of a variety of drug screening techniques. The peptide fragment employed in such a test may either be free in solution, affixed to a solid support, borne on a cell surface or located intracellularly. One may measure, for example, the formation of complexes between HJAK2 and the agent being tested. Alternatively, one can examine the diminution in complex formation between HJAK2 and a receptor caused by the agent being tested.

Methods of screening for drugs or any other agents which can affect macrophage activation comprise contacting such an agent with HJAK2 fragment and assaying for the presence of a complex between the agent and the HJAK2

1 fragment. In such assays, the HJAK2 fragment is typically labelled. After  
suitable incubation, free HJAK2 fragment is separated from that present in  
bound form, and the amount of free or uncomplexed label is a measure of the  
ability of the particular agent to bind to HJAK2.

6 Another technique for drug screening provides high throughput  
screening for compounds having suitable binding affinity to the HJAK2  
polypeptides and is described in detail in European Patent Application  
84/03564, published on September 13, 1984, incorporated herein by  
reference. Briefly stated, large numbers of different small peptide test  
11 compounds are synthesized on a solid substrate, such as plastic pins or  
some other surface. The peptide test compounds are reacted with HJAK2  
fragment and washed. Bound HJAK2 fragment is then detected by methods well  
known in the art. Purified HJAK2 can also be coated directly onto plates  
for use in the aforementioned drug screening techniques. In addition, non-  
neutralizing antibodies can be used to capture the peptide and immobilize  
16 it on the solid support.

21 This invention also contemplates the use of competitive drug  
screening assays in which neutralizing antibodies capable of binding HJAK2  
specifically compete with a test compound for binding to HJAK2 fragments.  
In this manner, the antibodies can be used to detect the presence of any  
peptide which shares one or more antigenic determinants with HJAK2.

#### XV Identification of Molecules Which Interact with HJAK2

26 The inventive purified HJAK2 is a research tool for identification,  
characterization and purification of interacting molecules. Appropriate  
labels are incorporated into HJAK2 by various methods known in the art and  
HJAK2 is used to capture soluble or interact with membrane-bound molecules.

31 A preferred method involves labeling the primary amino groups in HJAK2  
with <sup>125</sup>I Bolton-Hunter reagent (Bolton, AE and Hunter, WM (1973) Biochem J  
133: 529). This reagent has been used to label various molecules without  
concomitant loss of biological activity (Hebert CA et al (1991) J Biol Chem  
266: 18989-94; McColl S et al (1993) J Immunol 150:4550-4555). Membrane-  
bound molecules are incubated with the labelled HJAK2 molecules, washed to  
removed unbound molecules, and the HJAK2 complex is quantified. Data  
obtained using different concentrations of HJAK2 are used to calculate  
36 values for the number, affinity, and association of HJAK2.

Labelled HJAK2 fragments are also useful as a reagent for the

1 purification of molecules with which HJAK2 interacts, specifically  
including inhibitors. In one embodiment of affinity purification, HJAK2 is  
covalently coupled to a chromatography column. Cells and their membranes  
are extracted, HJAK2 is removed and various HJAK2-free subcomponents are  
passed over the column. Molecules bind to the column by virtue of their  
6 HJAK2 affinity. The HJAK2-complex is recovered from the column,  
dissociated and the recovered molecule is subjected to N-terminal protein  
sequencing or other identification procedure. If the captured molecule has  
an amino acid sequence, it can be used to design degenerate oligomers for  
use in cloning the gene from an appropriate cDNA library.

11 In an alternate method, monoclonal antibodies raised against HJAK2  
fragments are screened to identify those which inhibit the binding of  
labelled HJAK2. These monoclonal antibodies are then used in affinity  
purification or expression cloning of associated molecules. Other soluble  
binding molecules are identified in a similar manner. Labelled HJAK2 is  
16 incubated with extracts or other appropriate materials derived from lung,  
kidney or other tissues with activated monocytes or macrophages. After  
incubation, HJAK2 complexes (which are larger than the lone HJAK2 fragment)  
are identified by a sizing technique such as size exclusion chromatography  
or density gradient centrifugation and are purified by methods well known  
21 in the art. The soluble binding protein(s) are subjected to N-terminal  
sequencing to obtain information sufficient for database identification, if  
the soluble protein is known, or for cloning, if the soluble protein is  
unknown.

#### 26 XVI Use and Administration of Antibodies or Inhibitors to HJAK2

The antibodies and inhibitors can provide different effects when  
administered therapeutically. The antibodies and inhibitors are used to  
lessen or eliminate undue damage caused by disorders or diseases associated  
with upregulated HJAK2 expression. Each of these molecules or treatments  
31 (TSTs) will be formulated in a nontoxic, inert, pharmaceutically acceptable  
aqueous carrier medium preferably at a pH of about 5 to 8, more preferably  
6 to 8, although the pH may vary according to the different characteristics  
of the peptide, antibody or inhibitor being formulated and the condition to  
be treated. Characteristics of TSTs include solubility of the molecule,  
36 half-life, antigenicity/immunogenicity and the ability of the inhibitor to  
reach its target(s). These and other characteristics may aid in defining



1 an effective carrier. Native human proteins are preferred as TSTs, but  
recombinant peptides as well as organic or synthetic molecules resulting  
from drug screens may be equally effective in particular situations.

6 TSTs may be delivered by known routes of administration including but  
not limited to topical creams and gels; transmucosal spray and aerosol;  
transdermal patch and bandage; injectable, intravenous and lavage  
formulations; and orally administered liquids and pills particularly  
formulated to resist stomach acid and enzymes. The particular formulation,  
exact dosage, and route of administration will be determined by the  
attending physician and will vary according to each specific situation.

11 Such determinations are made by considering multiple variables such  
as the condition to be treated, the TST to be administered, and the  
pharmacokinetic profile of the particular TST. Additional factors which  
may be taken into account include disease state (eg. severity) of the  
patient, age, weight, gender, diet, time and frequency of administration,  
16 drug combination, reaction sensitivities, and tolerance/response to  
therapy. Long acting TST formulations might be administered every 3 to 4  
days, every week, or once every two weeks depending on half-life and  
clearance rate of the particular TST.

21 Normal dosage amounts may vary from 0.1 to 100,000 micrograms, up to  
a total dose of about 1 g, depending upon the route of administration.  
Guidance as to particular dosages and methods of delivery is provided in  
the literature. See US Patent No. 4,657,760; 5,206,344; or 5,225,212.  
Those skilled in the art will employ different formulations for different  
TSTs. Administration to lung cells may necessitate delivery in a manner  
26 different from that to kidney or other cells.

31 It is contemplated that conditions associated with altered HJAK2  
expression are treatable with TSTs. These conditions, which specifically  
include, but are not limited to, anemia, arteriosclerosis, asthma,  
bronchitis, emphysema, gingivitis, inflammatory bowel disease, insulin-  
dependent diabetes mellitus, leukemia, multiple endocrine neoplasias,  
osteoarthritis, osteoporosis, pulmonary fibrosis, rheumatoid arthritis,  
septic shock syndromes, and systemic lupus erythematosus may be  
specifically diagnosed by the tests discussed above. In addition, such  
tests may be used to monitor treatment.

1 All publications and patents mentioned in the above specification are  
herein incorporated by reference. Various modifications and variations of  
the described method and system of the invention will be apparent to those  
skilled in the art without departing from the scope and spirit of the  
invention. Although the invention has been described in connection with  
6 specific preferred embodiments, it should be understood that the invention  
as claimed should not be unduly limited to such specific embodiments.  
Indeed, various modifications of the above-described modes for carrying out  
the invention which are obvious to those skilled in the field of molecular  
biology or related fields are intended to be within the scope of the  
11 following claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Coleman, Roger  
Stuart, Susan G.

(ii) TITLE OF INVENTION: A NOVEL HUMAN JAK2 KINASE HOMOLOG

(iii) NUMBER OF SEQUENCES: 2

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: INCYTE PHARMACEUTICALS, INC.  
(B) STREET: 3174 Porter Drive  
(C) CITY: Palo Alto  
(D) STATE: CA  
(E) COUNTRY: USA  
(F) ZIP: 94304

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:  
(B) FILING DATE:  
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Luther, Barbara J.  
(B) REGISTRATION NUMBER: 33954  
(C) REFERENCE/DOCKET NUMBER: PF-0049 US

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 415-855-0555  
(B) TELEFAX: 415-852-0195

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 4482 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vii) IMMEDIATE SOURCE:

(A) LIBRARY: Placenta

(B) CLONE: 179527

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CCACGCGTCC GGTGCGCAAC CCGCAGGCGA CTGGGCGCTT CATCCCACCC TCACCCCTTT	60
CCAGCCAAGG TGGCTGATCG GAGTCAGGCT CTCGAGGTCG CATTGCCACG AAACGGNGTG	120
TGTGAGCGCG TTGTCCCCGG NCCCCGGGGC CACTTCCCCT CGGCCTAGNA GACTGGACTG	180
GGGAAGGACG GGTCTGTTGT ACCCGGGAGG TGGAAGGAAA AGCCGAAAGC GGAGAAGTGT	240
GCGGGAGGGG AGTCTCCGCG CGGAGGNAGA CCGGNCTCCT CCAGTGCAGG TTGTGCGCTG	300
GGGAGCCAGC CASGGCAAAT GTTCTGAAAA AGACTCTGCA TGGGAATGGC CTGCCTTACG	360
ATGACAGAAA TGGAGGGAAC ATCCACCTCT TCTATATATC AGAATGGTGA TATTTCTGGA	420
AATGCCAATT CTATGAAGCA AATAGATCCA GTTCTTCAGG TGTATCTTTA CCATTCCCTT	480
GGGAAATCTG AGGCAGATT TCTGACCTTT CCATCTGGGG AGTATGTTGG AGAAGAAATC	540
TGTATTGCTG CTTCTAAAGC TTGTGGTATC ACACCTGTGT ATCATAATAT GTTTGCTTTA	600
ATGAGTGAAA CAGAAAGGAT CTGGTATCCA CCCAACCATG TCTTCCATAT AGATGAGTCA	660
ACCAGGCATA ATGTACTCTA CAGAATAAGA TTTTACTTTC CTCGTTGGTA TTGCAGTGGC	720
AGCAACAGAG CCTATCGGCA TGGAATATCT CGAGGTGCTG AAGCTCCTCT TCTTGATGAC	780
TTTGTCATGT CTTACCTCTT TGCTCAGTGG CGGCATGATT TTGTGCATGG ATGGATAAAA	840
GTACCTGTGA CTCATGAAAC ACAGGAAGAA TGTCTTGGGA TGACAGTGTT AGATATGATG	900
AGAATAGCCA AAGAAAACGA TCAAACCCCA CTGGCCATCT ATAACTCTAT CAGCTACAAG	960
ACATTCTTAC CACAATGTAT TCGAGCAAAG ATCCAAGACT ATCATATTTT GACAAGGAAG	1020
CGAATAAGGT ACAGATTTTCG CAGATTTATT CAGCAATTCA GCCAATGCAA AGCCACTGCC	1080
AGAAACTTGA AACTTAAGTA TCTTATAAAT CTGGAAACTC TGCAGTCTGC CTTCTACACA	1140
GAGAAATTTG AAGTAAAAGA ACCTGGAAGT GGTCTTCAG GTGAGGAGAT TTTTGCAACC	1200
ATTATAATAA CTGGAAACGG TGGAATTCAG TGGTCAAGAG GGAAACATAA AGAAAGTGAG	1260

PF-0049 US

ACACTGACAG AACAGGATTT ACAGTTATAT TCGGATTTTC CTAATATTAT TGATGTCAGT	1320
ATTAAGCAAG CAAACCAAGA GGGTTCAAAT GAAAGCCGAG TTGTAAGTAT CCATAAGCAA	1380
GATGGTAAAA ATCTGGAAAT TGAAGTTAGC TCATTAAGGG AAGCTTTGTC TTTCGTGTCA	1440
TTAATTGATG GATATTATAG ATTAAGTACA GATGCACATC ATTACCTCTG TAAAGAAGTA	1500
GCACCTCCAG CCGTGCTTGA AAATATACAA AGCAACTGTC ATGGCCCAAT TTCGATGGAT	1560
TTTGCCATTA GTAAAGTGA GAAAGCAGGT AATCAGACTG GACTGTATGT ACTTCGATGC	1620
AGTCCTAAGG ACTTTAATAA ATATTTTTTG ACTTTTGCTG TCGAGCGAGA AAATGTCATT	1680
GAATATAAAC ACTGTTTGAT TACAAAAAAT GAGAATGAAG AGTACAACCT CAGTGGGACA	1740
AAGAAGAACT TCAGCAGTCT TAAAGATCTT TTGAATTGTT ACCAGATGGA AACTGTTTCGC	1800
TCAGACAATA TAATTTTCCA GTTTACTAAA TGCTGTCCCC CAAAGCCAAA AGATAAATCA	1860
AACCTTCTAG TCTTCAGAAC GAATGGTGTT TCTGATGTAC CAACCTCACC AACATTACAG	1920
AGGCCTACTC ATATGAACCA AATGGTGTTT CACAAAATCA GAAATGAAGA TTTGATATTT	1980
AATGAAAGCC TTGGCCAAGG CACTTTTACA AAGATTTTAA AAGGCGTACG AAGAGAAGTA	2040
GGAGACTACG GTCAACTGCA TGAAACAGAA GTTCTTTTAA AAGTTCTGGA TAAAGCACAC	2100
AGGAAGTATT CAGAGTCTTT CTTTGAAGCA GCAAGTATGA TGAGCAAGCT TTCTCACAAG	2160
CATTTGGTTT TAAATTATGG AGTATGTGTC TGTGGAGACG AGAATATTCT GGTTCAGGAG	2220
TTTGTAATAA TTGGATCACT AGATACATAT CTGAAAAAGA ATAAAAATTG TATAAATATA	2280
TTATGGAAAC TTGAAGTTGC TAAACAGTTG GCATGGGCCA TGCATTTTCT AGAAGAAAAC	2340
ACCCTTATTC ATGGGAATGT ATGTGCCAAA AATATTCTGC TTATCAGAGA AGAAGACAGG	2400
AAGACAGGAA ATCCTCCTTT CATCAAACCTT AGTGATCCTG GCATTAGTAT TACAGTTTTG	2460
CCAAAGGACA TTCTTCAGGA GAGAATACCA TGGGTACCAC CTGAATGCAT TGAAAATCCT	2520
AAAAATTTAA ATTTGGCAAC AGACAAATGG AGTTTTGGTA CCACTTTGTG GGAAATCTGC	2580
AGTGGAGGAG ATAAACCTCT AAGTGCTCTG GATTCTCAA GAAAGCTACA ATTTTATGAA	2640
GATAGGCATC AGCTTCCTGC ACCAAAGTGG GCAGAATTAG CAAACCTTAT AAATAATTGT	2700
ATGGATTATG AACCAGATTT CAGGCCTTCT TTCAGAGCCA TCATACGAGA TCTTAACAGT	2760

PF-0049 US

TTGTTTACTC CAGATTATGA ACTATTAACA GAAAATGACA TGTTACCAAA TATGAGGATA	2820
GGTGCCTTGG GGTTTTCTGG TGCCTTTGAA GACCGGGATC CTACACAGTT TGAAGAGAGA	2880
CATTTGAAAT TTCTACAGCA ACTTGGAAG GGTAATTTTG GGAGTGTGGA GATGTGCCGG	2940
TATGACCCTC TACAGGACAA CACTGGGGAG GTGGTCGCTG TAAAAAAGCT TCAGCATAGT	3000
ACTGAAGAGC ACCTAAGAGA CTTTGAAAGG GAAATTGAAA TCCTGAAATC CCTACAGCAT	3060
GACAACATTG TAAAGTACAA GGGAGTGTGC TACAGTGCTG GTCGGCGTAA TCTAAAATTA	3120
ATTATGGAAT ATTTACCATA TGGAAGTTTA CGAGACTATC TTCAAAAACA TAAAGAACGG	3180
ATAGATCACA TAAAACTTCT GCAGTACACA TCTCAGATAT GCAAGGGTAT GGAGTATCTT	3240
GGTACAAAAA GGTATATCCA CAGGGATCTG GCAACGAGAA ATATATTGGT GGAGAACGAG	3300
AACAGAGTTA AAATTGGRGA TTTTGGGTTA ACCAAAGTCT TGCCACAAGA CAAAGAATAC	3360
TATAAAGTAA AAGAACCTGG TGAAAGTCCC ATATTCTGGT ATGCTCCAGA ATCACTGACA	3420
GAGAGCAAGT TTTCTGTGGC CTCAGATGTT TGGAGCTTTG GAGTGGTTCT GTATGAACTT	3480
TTCACATACA TTGAGAAGAG TAAAAGTCCA CCAGCGGAAT TTATGCGTAT GATTGGCAAT	3540
GACAAACAAG GACAGATGAT CGTGTTCCAT TTGATAGAAC TTTTGAAGAA TAATGGAAGA	3600
TTACCAAGAC CAGATGGATG CCCAGATGAG ATCTATATGA TCATGACAGA ATGCTGGAAC	3660
AATAATGTAA ATCAACGCCC CTCCTTTAGG GATCTAGCTC TTCGAGTGGA TCAAATAAGG	3720
GATAACATGG CTGGATGAAA GAAATGACCT TCATTCTGAG ACCAAAGTAG ATTTACAGAA	3780
CAAAGTTTTA TATTTACAT TGCTGTGGAC TATTATTACA TATATCATT TATATAAAT	3840
CATGATGCTA GCCAGCAAAG ATGTGAAAAT ATCTGCTCAA AACTTTCAA GTTTAGTAAG	3900
TTTTTCTTCA TGAGGCCACC AGTAAAAGAC ATTAATGAGA ATTCCTTAGC AAGGATTTTG	3960
TAAGAAGTTT CTTAAACATT GTCAGTTAAC ATCACTCTTG TCTGGCAAAA GAAAAAAAT	4020
AGACTTTTTT AACTCAGCTT TTTGAGACCT GAAARAATTA TTATGTAAAT TTTGCAATGT	4080
TAAAGATGCA CAGAATATGT ATGTATAGTT TTTACCACAG TGGATGTATA ATACCTGGC	4140
ATCTTGTGTG ATGTTTACA CACATGAGGG CTGGTGTTC TTAATACTGT TTTCTAATTT	4200
TTCCATGGTT AATCTATAAT TAATTACTTC ACTAAACAAA CAAATTAAGA TGTTCAGATA	4260

PF-0049 US

ATTGAATAAG TACCTTTGTG TCCTTGTTCA TTTATATCGC TGGCCAGCAT TATAAGCAGG 4320  
TGTATACTTT TAGCTTGTAG TTCCATGTAC TGTAATATT TTTCACATAA AGGGAACAAA 4380  
TGTCTAGTTT TATTTGTATA GGAAATTTGC CCTGACCCTA AATAATACAT TTTGAAATGA 4440  
AACAAAGCTTA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AG 4482

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1132 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Gly Met Ala Cys Leu Thr Met Thr Glu Met Glu Gly Thr Ser Thr  
1 5 10 15  
Ser Ser Ile Tyr Gln Asn Gly Asp Ile Ser Gly Asn Ala Asn Ser Met  
20 25 30  
Lys Gln Ile Asp Pro Val Leu Gln Val Tyr Leu Tyr His Ser Leu Gly  
35 40 45  
Lys Ser Glu Ala Asp Tyr Leu Thr Phe Pro Ser Gly Glu Tyr Val Gly  
50 55 60  
Glu Glu Ile Cys Ile Ala Ala Ser Lys Ala Cys Gly Ile Thr Pro Val  
65 70 75 80  
Tyr His Asn Met Phe Ala Leu Met Ser Glu Thr Glu Arg Ile Trp Tyr  
85 90 95  
Pro Pro Asn His Val Phe His Ile Asp Glu Ser Thr Arg His Asn Val  
100 105 110  
Leu Tyr Arg Ile Arg Phe Tyr Phe Pro Arg Trp Tyr Cys Ser Gly Ser  
115 120 125  
Asn Arg Ala Tyr Arg His Gly Ile Ser Arg Gly Ala Glu Ala Pro Leu  
130 135 140  
Leu Asp Asp Phe Val Met Ser Tyr Leu Phe Ala Gln Trp Arg His Asp  
145 150 155 160

PF-0049 US

Phe Val His Gly Trp Ile Lys Val Pro Val Thr His Glu Thr Gln Glu  
165 170 175

Glu Cys Leu Gly Met Thr Val Leu Asp Met Met Arg Ile Ala Lys Glu  
180 185 190

Asn Asp Gln Thr Pro Leu Ala Ile Tyr Asn Ser Ile Ser Tyr Lys Thr  
195 200 205

Phe Leu Pro Gln Cys Ile Arg Ala Lys Ile Gln Asp Tyr His Ile Leu  
210 215 220

Thr Arg Lys Arg Ile Arg Tyr Arg Phe Arg Arg Phe Ile Gln Gln Phe  
225 230 235 240

Ser Gln Cys Lys Ala Thr Ala Arg Asn Leu Lys Leu Lys Tyr Leu Ile  
245 250 255

Asn Leu Glu Thr Leu Gln Ser Ala Phe Tyr Thr Glu Lys Phe Glu Val  
260 265 270

Lys Glu Pro Gly Ser Gly Pro Ser Gly Glu Glu Ile Phe Ala Thr Ile  
275 280 285

Ile Ile Thr Gly Asn Gly Gly Ile Gln Trp Ser Arg Gly Lys His Lys  
290 295 300

Glu Ser Glu Thr Leu Thr Glu Gln Asp Leu Gln Leu Tyr Cys Asp Phe  
305 310 315 320

Pro Asn Ile Ile Asp Val Ser Ile Lys Gln Ala Asn Gln Glu Gly Ser  
325 330 335

Asn Glu Ser Arg Val Val Thr Ile His Lys Gln Asp Gly Lys Asn Leu  
340 345 350

Glu Ile Glu Leu Ser Ser Leu Arg Glu Ala Leu Ser Phe Val Ser Leu  
355 360 365

Ile Asp Gly Tyr Tyr Arg Leu Thr Ala Asp Ala His His Tyr Leu Cys  
370 375 380

Lys Glu Val Ala Pro Pro Ala Val Leu Glu Asn Ile Gln Ser Asn Cys  
385 390 395 400

His Gly Pro Ile Ser Met Asp Phe Ala Ile Ser Lys Leu Lys Lys Ala  
405 410 415

Gly Asn Gln Thr Gly Leu Tyr Val Leu Arg Cys Ser Pro Lys Asp Phe  
420 425 430



PF-0049 US

Asn Lys Tyr Phe Leu Thr Phe Ala Val Glu Arg Glu Asn Val Ile Glu  
435 440 445

Tyr Lys His Cys Leu Ile Thr Lys Asn Glu Asn Glu Glu Tyr Asn Leu  
450 455 460

Ser Gly Thr Lys Lys Asn Phe Ser Ser Leu Lys Asp Leu Leu Asn Cys  
465 470 475 480

Tyr Gln Met Glu Thr Val Arg Ser Asp Asn Ile Ile Phe Gln Phe Thr  
485 490 495

Lys Cys Cys Pro Pro Lys Pro Lys Asp Lys Ser Asn Leu Leu Val Phe  
500 505 510

Arg Thr Asn Gly Val Ser Asp Val Pro Thr Ser Pro Thr Leu Gln Arg  
515 520 525

Pro Thr His Met Asn Gln Met Val Phe His Lys Ile Arg Asn Glu Asp  
530 535 540

Leu Ile Phe Asn Glu Ser Leu Gly Gln Gly Thr Phe Thr Lys Ile Phe  
545 550 555 560

Lys Gly Val Arg Arg Glu Val Gly Asp Tyr Gly Gln Leu His Glu Thr  
565 570 575

Glu Val Leu Leu Lys Val Leu Asp Lys Ala His Arg Asn Tyr Ser Glu  
580 585 590

Ser Phe Phe Glu Ala Ala Ser Met Met Ser Lys Leu Ser His Lys His  
595 600 605

Leu Val Leu Asn Tyr Gly Val Cys Val Cys Gly Asp Glu Asn Ile Leu  
610 615 620

Val Gln Glu Phe Val Lys Phe Gly Ser Leu Asp Thr Tyr Leu Lys Lys  
625 630 635 640

Asn Lys Asn Cys Ile Asn Ile Leu Trp Lys Leu Glu Val Ala Lys Gln  
645 650 655

Leu Ala Trp Ala Met His Phe Leu Glu Glu Asn Thr Leu Ile His Gly  
660 665 670

Asn Val Cys Ala Lys Asn Ile Leu Leu Ile Arg Glu Glu Asp Arg Lys  
675 680 685

Thr Gly Asn Pro Pro Phe Ile Lys Leu Ser Asp Pro Gly Ile Ser Ile  
690 695 700

PF-0049 US

Thr Val Leu Pro Lys Asp Ile Leu Gln Glu Arg Ile Pro Trp Val Pro  
705 710 715 720

Pro Glu Cys Ile Glu Asn Pro Lys Asn Leu Asn Leu Ala Thr Asp Lys  
725 730 735

Trp Ser Phe Gly Thr Thr Leu Trp Glu Ile Cys Ser Gly Gly Asp Lys  
740 745 750

Pro Leu Ser Ala Leu Asp Ser Gln Arg Lys Leu Gln Phe Tyr Glu Asp  
755 760 765

Arg His Gln Leu Pro Ala Pro Lys Trp Ala Glu Leu Ala Asn Leu Ile  
770 775 780

Asn Asn Cys Met Asp Tyr Glu Pro Asp Phe Arg Pro Ser Phe Arg Ala  
785 790 795 800

Ile Ile Arg Asp Leu Asn Ser Leu Phe Thr Pro Asp Tyr Glu Leu Leu  
805 810 815

Thr Glu Asn Asp Met Leu Pro Asn Met Arg Ile Gly Ala Leu Gly Phe  
820 825 830

Ser Gly Ala Phe Glu Asp Arg Asp Pro Thr Gln Phe Glu Glu Arg His  
835 840 845

Leu Lys Phe Leu Gln Gln Leu Gly Lys Gly Asn Phe Gly Ser Val Glu  
850 855 860

Met Cys Arg Tyr Asp Pro Leu Gln Asp Asn Thr Gly Glu Val Val Ala  
865 870 875 880

Val Lys Lys Leu Gln His Ser Thr Glu Glu His Leu Arg Asp Phe Glu  
885 890 895

Arg Glu Ile Glu Ile Leu Lys Ser Leu Gln His Asp Asn Ile Val Lys  
900 905 910

Tyr Lys Gly Val Cys Tyr Ser Ala Gly Arg Arg Asn Leu Lys Leu Ile  
915 920 925

Met Glu Tyr Leu Pro Tyr Gly Ser Leu Arg Asp Tyr Leu Gln Lys His  
930 935 940

Lys Glu Arg Ile Asp His Ile Lys Leu Leu Gln Tyr Thr Ser Gln Ile  
945 950 955 960

Cys Lys Gly Met Glu Tyr Leu Gly Thr Lys Arg Tyr Ile His Arg Asp  
965 970 975

PF-0049 US

Leu Ala Thr Arg Asn Ile Leu Val Glu Asn Glu Asn Arg Val Lys Ile  
980 985 990

Gly Ile Leu Gly \* Pro Lys Ser Cys His Lys Thr Lys Asn Thr Ile  
995 1000 1005

Lys \* Lys Asn Leu Val Lys Val Pro Tyr Ser Gly Met Leu Gln Asn  
1010 1015 1020

His \* Gln Arg Ala Ser Phe Leu Trp Pro Gln Met Phe Gly Ala Leu  
1025 1030 1035 1040

Glu Trp Phe Cys Met Asn Phe Ser His Thr Leu Arg Arg Val Lys Val  
1045 1050 1055

His Gln Arg Asn Leu Cys Val \* Leu Ala Met Thr Asn Lys Asp Arg  
1060 1065 1070

\* Ser Cys Ser Ile \* \* Asn Phe \* Arg Ile Met Glu Asp Tyr  
1075 1080 1085

Gln Asp Gln Met Asp Ala Gln Met Arg Ser Ile \* Ser \* Gln Asn  
1090 1095 1100

Ala Gly Thr Ile Met \* Ile Asn Ala Pro Pro Leu Gly Ile \* Leu  
1105 1110 1115 1120

Phe Glu Trp Ile Lys \* Gly Ile Thr Trp Leu Asp  
1125 1130